

## 17 Arm- en handfunctiestoornissen

### Literatuur

- Alon G, McBride K. Persons with C5 or C6 tetraplegia achieve selected functional gains using a neuroprosthesis. *Arch Phys Med Rehabil* 2003; 84: 119-24.
- Bedbrook G. *The care and management of spinal cord injury*. New York-Heidelberg-Berlin: Springer Verlag; 1981.
- Brucker BS, Bulaeveva NV. Biofeedback effect on electromyography responses in patients with spinal cord injury. *Arch Phys Med Rehabil* 1996; 77: 133-37.
- Bryden AM, Wuolle KS, Murray PK, Peckham PH. Perceived outcomes of upper extremity surgical reconstruction in individuals with tetraplegia at model spinal cord injury systems. *Spinal Cord* 2004; 42: 169-76.
- Catz A, Itzkovich M, Agranov E et al. SCIM-Spinal cord independence measure. A new disability scale for patients with spinal cord lesions. *Spinal Cord* 1997; 35: 850-6.
- Chea J, Triolo RJ, Kilgore KL, Creasey G, Dimarco AF. Neuromuscular electrical stimulation in spinal cord injury. In: Kirschblum S, Campagnolo DI, DeLisa JA, editors. *Spinal Cord Medicine*. Philadelphia: Lippincott, Williams & Wilkins; 2002. Hoofdstuk 25.
- Chen D, Jaeger RJ. Functional electrical stimulation. Technical advances and clinical applications. *Physical Medicine and Rehabilitation* 1997; 11: 39-53.
- Coulet B, Allieu Y, Chammas M et al. *Variability of the spinal cord injury zone extent in high level tetraplegia and consequences on conventional functional surgery indications especially in key grip construction*. Abstract book 7th International Conference on Tetraplegia: Surgery and Rehabilitation; June 2001 Bologna, Italy, p. 25.
- Curtin M. Development of a tetraplegic hand assessment and splinting protocol. *Paraplegia* 1994; 32: 159-69.
- Ditunno JF, Flanders AE, Kirshblum S et al. Predicting outcome in traumatic spinal cord injury. In: Kirshblum S, Campagnolo DI, DeLisa JA, editors. *Spinal Cord Medicine*. Philadelphia: Lippincott, Williams & Wilkins; 2002.
- Doll U, Maurer B, Spahn B, Fromm B. Functional hand development in tetraplegia. *Spinal Cord* 1998; 36: 818-21.
- Forner-Cordero I, Mudarra-Garcia J, Forner-Valero JV, Vilar-de-la-Pena R. The role of upper limb surgery in tetraplegia. *Spinal Cord* 2003; 41: 90-6.
- Franceschini M, Di Clemente B, Rampello A et al. Longitudinal outcome 6 years after spinal cord injury. *Spinal Cord* 2003; 41: 280-5.
- Gorman PH, Hunter Peckham P. Upper extremity functional neuromuscular stimulation. *Neuro Rehab* 1991; 5: 3-11.
- Handa Y, Hoshiyama N. Functional electrical stimulation for the control of the upper extremities. *Medical Progress through Technology* 1987; 12: 51-63.

## Handboek dwarslaesierevalidatie

- Handa Y, Ohkubo K, Hoshiyama N. A portable multichannel FES-system for restoration of upper motor function of the paralyzed extremities. *Automedica* 1989; 11: 221.
- Hartkopp A. *Training of wrist extensor muscles by electrical stimulation in spinal cord injured tetraplegic individuals with special reference to contractile and metabolic properties* [dissertatie]. University of Copenhagen; April 1999.
- Harvey L. Principles of conservative management for a non-orthotic tenodesis grip in tetraplegics. *J Hand Ther* 1996; 9: 238-42.
- Harvey LA, Batty J, Jones R, Crosbie J. Hand function of C6 and C7 tetraplegics 1-16 years following injury. *Spinal Cord* 2001; 39: 37-43.
- Hentz VR, Leclercq C. *Surgical rehabilitation of the upper limb in tetraplegia*. London-Edinburgh-New York-Philadelphia-St Louis-Sydney-Toronto: WB Saunders; 2002.
- House JH, Walsh T. Two stage reconstruction of the tetraplegic hand. In: Strickland JW, editor. *Master Techniques in Orthopedic Surgery, the Hand*. Philadelphia: Lippincott-Raven; 2002.
- Hummel MJ, Snoek GJ, Til JA van, Rossum W van, IJzerman MJ. A multicriteria decision analysis of augmentative treatment of upper limbs in persons with tetraplegia. *J Rehabil Res Dev* 2005; 42: 635-44, online publication [www.vard.org/jour/05/42/5/Hummel.html](http://www.vard.org/jour/05/42/5/Hummel.html).
- Johnstone BR, Jordan CJ, App B, Buntine JA. A Review of Surgical Rehabilitation of the Upper Limb in Quadriplegia. *Paraplegia* 1988; 26: 317-39.
- Keith MW, Kilgore KI, Hunter Peckham P et al. Tendon transfer and functional electrical stimulation for restoration of hand function in spinal cord injury. *J Hand Surg* 1996; 21A: 89-99.
- Keith MW, Lacey H. Surgical rehabilitation of the tetraplegic upper extremity. *J Neurol Rehabil* 1991; 5: 75-87.
- Kirshblum S, Campagnolo DI, DeLisa JA, editors. *Spinal Cord Medicine*. Philadelphia: Lippincott, Williams & Wilkins; 2002. Hoofdstuk 19: pp 275-282; hoofdstuk 25: pp 360-7; hoofdstuk 28.
- Kirshblum S, Ho CH, House JG, Druin E, Nead C, Drastal S. Rehabilitation of Spinal Cord Injury. In: Kirshblum S, Campagnolo DI, DeLisa JA, editors. *Spinal Cord Medicine*. Philadelphia: Lippincott, Williams & Wilkins; 2002. Hoofdstuk 19.
- Kohlmeyer KM, Hill JP, Yarkony GM, Jaeger RJ. Electrical stimulation and biofeedback on recovery of tenodesis grasp. A controlled study. *Arch Phys Med Rehabil* 1996; 77: 702-6.
- Lamb, DW, Chan KM. Surgical reconstruction of the upper limb in traumatic quadriplegia. *J Bone Joint Surg Am* 1983; 65B: 291-8.
- Long C, Lawton EB. Functional significance of spinal cord injury level. *Arch Phys Med Rehabil* 1955; 36: 249-55.
- Marino R, Goin J. Development of a short form quadriplegia index of function scale. *Spinal Cord* 1999; 37: 289-96.

## Handboek dwarslaesierevalidatie

- Marino RJ, Huang M, Knight P et al. Assessing selfcare status in quadriplegia. Comparison of the Quadriplegia Index of Function (QIF) and the Functional Independence Measure (FIM). *Paraplegia* 1993; 31: 225-33.
- Marino RJ, Rider Foster D, Maissel G, Ditunno JF. Superiority of motor level over single neurological level in categorising tetraplegia. *Paraplegia* 1993; 31: 510-13.
- Marino RJ, Stineman MG. Functional assessment in spinal cord injury. *Top Spinal Cord Inj Rehabil* 1996; 1: 32-45.
- Mc Dowell CL, Moberg M, House JH. The Second International Conference on Surgical Rehabilitation of the Upper Limb in Tetraplegia. *J Hand Surg* 1986; 11A: 604-8.
- Mc Dowell CL. Tetraplegia. In: Green DP, editor. *Operative hand surgery*. 3rd ed. New York-Edinburgh-London-Melbourne-Tokyo: Churchill Livingstone.
- Meiners T, Abel R, Lindel K, Mesecke U. Improvements in activities of daily living following functional hand surgery for treatment of lesions to the cervical spinal cord. Self-assessment by patients. *Spinal Cord* 2002; 40: 574-80.
- Moberg E. *The upper limb in tetraplegia*. Stuttgart: Georg Thieme Verlag; 1978.
- Mulcahey MJ, Betz RR, Kozin SH, Smith BT, Hutchinson D, Lutz C. Implantation of the Freehand System during initial rehabilitation using minimally invasive techniques. *Spinal Cord* 2004; 42: 146-55.
- Murphy C, Chuinard R. Management of upper extremity in traumatic tetraplegia. *Hand Clinics* 1998; 4: 201-9.
- Nathan RH. FNS of the upper limb. Targeting the forearm muscles for surface stimulation. *Medical and Biological Engineering and Computing* 1990; 28: 249-56.
- Peckham PH, Keith MW, Kilgore KL et al. Efficacy of an implanted neuroprosthesis for restoring hand grasp in tetraplegia. A multicenter study. *Arch Phys Med Rehabil* 2001; 82: 1380-88.
- Peckham PH, Kilgore L, Keith MW et al. An advanced neuroprosthesis for restoration of hand and upper arm control using an implantable controller. *J Hand Surg* 2002; 27A: 265-75.
- Pons C, Post M, Lieshout G van. The Van Lieshout Test for Hand function. *J Rehabil Res Dev* 2004; 41 Suppl 2: 61.
- Popovic D, Stojanovic A, Pjanovic A et al. Clinical evaluation of the Bionic Glove. *Arch Phys Med Rehabil* 1999; 80: 299-304.
- Popovic MB, Popovic DB, Sinkjaer T et al. Clinical evaluation of functional electrical therapy in hemiplegia. *J Rehabil Res Dev* 2003; 40: 443-54.
- Popovic MR, Thrasher TA, Adams ME et al. Functional electrical therapy. Retraining grasping in spinal cord injury. *Spinal Cord* 2006; 44: 143-51.
- Post M. *Living with Spinal Cord Injury* [dissertatie]. University of Utrecht: 22-4-1997.
- Post MWM, Lieshout G van, Snoek GJ et al. Validity of the Van Lieshout Test for Hand Function in Tetraplegia. *J Rehabil Res Dev* 2004; 41 Suppl 2: 62.

## Handboek dwarslaesierevalidatie

- Prochazka A, Gauthier M, Wieler M et al. The Bionic Glove. An electrical stimulator garment that provides controlled grasp and hand opening in quadriplegia. *Arch Phys Med Rehabil* 1997; 78: 608-14.
- Rothwell AG, Sinnott A, Mohammed KD et al. Upper limb surgery for tetraplegia. A 10 year re-review of hand function. *J Hand Surg* 2003; 28A: 489-95.
- Snoek GJ, IJzerman MJ, Groen FAC in 't et al. Use of de NESS Handmaster to restore handfunction in tetraplegia. Clinical experiences in 10 patients. *Spinal Cord* 2000; 38: 244-9.
- Snoek GJ. *Patient preferences for reconstructive interventions of the upper limb in tetraplegia* [dissertatie]. University of Twente: 8-12-2005.
- Stroh-Wuolle K, Doren CL van, Thorpe GB et al. Development of a Quantitative hand grasp and release test for patients with tetraplegia using a hand neuroprosthesis. *J Hand Surg* 1994; 19A: 209-18.
- Taylor P, Esnouf J, Hobby J. Pattern of use and user satisfaction of Neuro Control freehand system. *Spinal Cord* 2001; 39: 156-60.
- Triolo R, Nathan R, Handa Y et al. Challenges to clinical deployment of upper limb neuroprostheses. *J Rehab Res Dev* 1996; 33: 111-22.
- Tuijl JH van, Janssen-Potten YJM, Seelen HAM. Evaluation of upper extremity motor function tests in tetraplegia. *Spinal Cord* 2002; 40: 51-64.
- Waters R, Sie JH, Gellman H, Tognella M. Functional hand surgery following tetraplegia. *Arch Phys Med Rehabil* 1996; 77: 86-94.
- Waters RL, Mucitelli LM. Tendon transfer to improve function of patients with tetraplegia. In: Kirshblum S, Campagnolo DI, DeLisa JA, editors. *Spinal Cord Medicine*. Philadelphia: Lippincott, Williams & Wilkins; 2002. Hoofdstuk 28.
- Wuolle KS, Bryden AM, Peckham PH, Murray PK, Keith M. Satisfaction with upper extremity surgery in individuals with tetraplegia. *Arch Phys Med Rehabil* 2003; 84: 1145-49.
- Yarkony GM, Roth EJ, Cybulski G et al. Neuromuscular stimulation in spinal cord injury. Restoration of functional movement of the extremities. *Arch Phys Med Rehabil* 1992; 73: 78-83.
- Yavuz N, Tezyurek M, Akyuz M. A comparison of two functional tests in quadriplegia. The quadriplegia index of function and the functional independence measure. *Spinal Cord* 1998; 36: 832-7.
- Zalfonte RD, Demagone DA, Herbison GJ, Ditunno JF. Daily self care in quadriplegic subjects. *Neurol Rehabil* 1991; 4: 17-24.
- Zancolli EA. Midcervical tetraplegia with strong wrist extension: a two-stage synergistic reconstruction of the hand. *Hand Clinics* 2002; 18: 481-95.
- Zancolli E. Surgery for the quadriplegic hand with active strong wrist extension preserved. A study of 97 cases. *Clinical Orthopaedics* 1975; 112: 102-13.